

Provided for non-commercial research and education use.
Not for reproduction, distribution or commercial use.



This article appeared in a journal published by Elsevier. The attached copy is furnished to the author for internal non-commercial research and education use, including for instruction at the authors institution and sharing with colleagues.

Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited.

In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit:

<http://www.elsevier.com/authorsrights>



Contents lists available at SciVerse ScienceDirect

Journal of Human Evolution

journal homepage: www.elsevier.com/locate/jhevol

The oldest human fossil in Europe, from Orce (Spain)



Isidro Toro-Moyano^{a,*}, Bienvenido Martínez-Navarro^{b,**}, Jordi Agustí^b,
 Caroline Souday^{c,d}, José María Bermúdez de Castro^e, María Martínón-Torres^e,
 Beatriz Fajardo^{f,g}, Mathieu Duval^{d,e}, Christophe Falguères^d, Oriol Oms^h,
 Josep Maria Parés^e, Pere Anadónⁱ, Ramón Juliàⁱ, José Manuel García-Aguilar^j,
 Anne-Marie Moigne^{d,k}, María Patrocínio Espigares^l, Sergio Ros-Montoya^l,
 Paul Palmqvist^m

^a Museo Arqueológico de Granada, Carrera del Darro 41-43, 18010 Granada, Spain

^b ICREA, Institut de Paleoeecologia Humana i Evolució Social, Àrea de Prehistoria, Universitat Rovira i Virgili, C/Marcel·lí Domingo s/n, Campus Sescelades, 43007 Tarragona, Spain

^c Center for the Study of Human Origins – CSHO, 25 Waverly Place, New York, NY 10003-6790, USA

^d Muséum National d'Histoire Naturelle, Département de Préhistoire, UMR 7194 du CNRS, 1 rue René Panhard, 75013 Paris, France

^e Centro Nacional de Investigación sobre la Evolución Humana (CENIEH), Paseo de Atapuerca, 09002 Burgos, Spain

^f Université Paul Valéry, Montpellier III, UMR 5140 «Archéologie des sociétés méditerranéennes» Route de Mende, F-34199 Montpellier, France

^g Stone Age Institute, Indiana University, 1392 W. Ditterrore Rd., Gasport, IN 47433, USA

^h Departament de Geologia, Universitat Autònoma de Barcelona, Bellaterra, Spain

ⁱ Institut de Ciències de la Terra 'J. Almera' (CSIC), C/L. Solé Sabarís sn, 08028 Barcelona, Spain

^j Departamento de Estratigrafía y Paleontología, Campus de Fuentenueva, Universidad de Granada, 18003 Granada, Spain

^k Centre européen de recherches préhistoriques, avenue Léon-Grégoire, 66720 Tautavel, France

^l Museo de Prehistoria y Paleontología, Orce, Granada 18858, Spain

^m Departamento de Geología y Ecología, Universidad de Málaga, Campus de Teatinos, 29071 Málaga, Spain

ARTICLE INFO

Article history:

Received 8 October 2011

Accepted 21 January 2013

Available online 05 March 2013

Keywords:

Human tooth

Early Pleistocene

Barranco León

ABSTRACT

The Orce region has one of the best late Pliocene and early Pleistocene continental paleobiological records of Europe. It is situated in the northeastern sector of the intramontane Guadix-Baza Basin (Granada, Andalusia, southern Spain). Here we describe a new fossil hominin tooth from the site of Barranco León, dated between 1.02 and 1.73 Ma (millions of years ago) by Electron Spin Resonance (ESR), which, in combination with paleomagnetic and biochronologic data, is estimated to be close to 1.4 Ma. While the range of dates obtained from these various methods overlaps with those published for the Sima del Elefante hominin locality (1.2 Ma), the overwhelming majority of evidence points to an older age. Thus, at the moment, the Barranco León hominin is the oldest from Western Europe.

© 2013 Elsevier Ltd. All rights reserved.

Introduction

The date of the earliest human occupation of Europe has been recently challenged by new paleontological and archaeological

evidence from both longitudinal extremes of the continent. The Georgian site of Dmanisi has provided an important collection of Oldowan lithic artefacts and very significant fossils of early *Homo*, ca. 1.8 Ma (millions of years ago) (Vekua et al., 2002; Lordkipanidze et al., 2007; Ferring et al., 2011), the Spanish site of Sima del Elefante in Atapuerca has yielded some lithic artefacts and a mandibular fragment dated to ~1.2 Ma (Carbonell et al., 2008), the Italian site of Pirro Nord (1.3–1.7 Ma) (Arzarello et al., 2009), and the French sites of Lézignan-la-Cèbe and Pont-de-Lavaud (1.57 and ~1.1 Ma, respectively) (Crochet et al., 2009; Despriée et al., 2010) have provided some lithic artefacts, and two early Pleistocene localities placed in the northeastern sector of the Guadix-Baza basin in Spain, Fuente Nueva-3 and Barranco León, have yielded huge lithic assemblages and evidence of anthropic use of tools on large mammal bones (Turq et al., 1996; Martínez-Navarro et al., 1997;

* Corresponding author.

** Corresponding author.

E-mail addresses: isidro.toro@juntadeandalucia.es (I. Toro-Moyano), bienvenido.martinez@icrea.cat (B. Martínez-Navarro), jordi.agusti@icrea.cat (J. Agustí), caroline.souday@gmail.com (C. Souday), jm.ber@cenieh.es (J.M. Bermúdez de Castro), maria.martinon@cenieh.es (M. Martínón-Torres), bea.fajardo@gmail.com (B. Fajardo), mathieu.duval@cenieh.es (M. Duval), falguere@mnhn.fr (C. Falguères), joseporiol.oms@uab.cat (O. Oms), josep.pares@cenieh.es (J.M. Parés), panadon@ija.csic.es (P. Anadón), ramon.julia@ija.csic.es (R. Julià), jmg.aguilar@gmail.com (J.M. García-Aguilar), ammoigne@tautavel.univ-perp.fr (A.-M. Moigne), mpepigares@gmail.com (M.P. Espigares), sergiorosm@gmail.com (S. Ros-Montoya), ppb@uma.es (P. Palmqvist).

Gibert et al., 1998b; Toro-Moyano et al., 2003, 2009, 2011; Palmqvist et al., 2005; Espigares et al., 2013).

Here we report the finding of a first deciduous molar of early *Homo* dating back to an age older than 1.2 Ma, probably close to 1.4 Ma, from Orce, which was found in direct association with assemblages of lithic artefacts and large mammal bones (Oms et al., 2000b; Toro-Moyano et al., 2009; Martínez-Navarro et al., 2010).

The tooth comes from level BL D (previously referred to as BL 5; i.e., Martínez-Navarro et al., 2005), square J54, of the section at Barranco León, a tributary creek of the Orce river, in the northeast of the Plio-Pleistocene lacustrine deposits of the Guadix-Baza Basin (Vera, 1970; Sanz de Galdeano and Vera, 1992; Fernández et al., 1996) (Fig. 1). The Barranco León section spans the middle terrigenous member (lowest part of section in Fig. 1) and the upper 'Silty Calcareous Member' of the Baza Formation (Vera et al., 1985; Oms et al., 2000a, 2011), which is dominated mainly by limestones, sandstones, carbonate silts and dark mudstones, deposited in a lake with an alternation of slightly saline and saline waters (Anadón et al., 1994; Anadón and Gabàs, 2009). The excavated layers of the BL section show sediments associated with a swampy environment, except level D, which shows fluvial features and encloses the archaeological level (see further details in Oms et al., 2011). Level BL D contains abundant remains of early Pleistocene large and small mammals composed of *Ursus* sp., *Canis mosbachensis*, *Lycaon lycaonoides*, *Vulpes* cf. *praeglacialis*, *Pachycrocuta brevirostris*, *Meles* sp., *Stephanorhinus hundsheimensis*, *Equus altidens*, *Equus sussenbornensis*, *Hippopotamus antiquus*, *Bison* sp., *Hemitragus* cf. *albus*, *Praemegaceros verticornis*, *Metacervoceros rhenanus*, *Oryctolagus* cf. *lacosti*, *Mimomys savini*, *Allophaiomys* aff. *lavocati*, *Allophaiomys* sp., *Apodemus* aff. *mystacinus*, *Histrix* sp., *Galemys* sp., *Erinacena* indet., *Crociodura* sp., *Sorex minutus*, and *Sorex* sp. (Agustí and Madurell, 2003; Furió-Bruno, 2003; Martínez-Navarro et al., 2010 and references there in), and stone artefacts (Oms et al., 2000b) that are broadly similar to other African and European lithic assemblages assigned to the Oldowan Industry Complex, or Mode I technology, characterized by a low degree of standardization (i.e., simple core forms and debitage).

Although some controversial 'hypothetical human remains' from the Orce site of Venta Micena were published during the last three decades (Gibert et al., 1983, 1998a, 2002; Campillo et al., 2006; among others), including cranial fragment VM-0 and limb bone diaphyses VM-1960, VM-3691 and VM-12000, none of these specimens has enough anatomical resolution to be ascribed to the genus *Homo* and they were refuted by Moyà-Solà and Köhler (1997), Palmqvist (1997), Martínez-Navarro (2002), and Palmqvist et al. (2005), among others. Finally, there is also a tooth fragment (BL5-0) found out of stratigraphic context in 1994 during a geological survey at the Barranco León ravine by A. Arribas. This specimen, which was not unearthed from the excavation quarry of the site, was preliminarily identified as a putative hominin fossil by Arribas. It is a small tooth fragment, which preserves the mesial part of the crown and root (4.6 mm crown height, 2.9 mm root length). The tooth has an advanced degree of wearing, showing the exposed dentine and a reduced pulp cavity. Following the comparative study of the striae of Retzius and Hunter-Schreger bands, the shape of the enamel prism patterns and the enamel thickness (which is clearly thinner than in human teeth), Gibert et al. (1999) published the tooth fragment as part of an upper human molar of early Pleistocene age. However, such conclusion should not be accepted without caution, given the reservations on the stratigraphic position of the finding (Arribas and Palmqvist, 2002) and the possibility that it could belong to a deciduous tooth of *H. antiquus*, one of the large mammal species that are better represented in the faunal assemblage preserved in the swampy-lacustrine facies of Barranco León (Palmqvist et al., 2005). More specifically, the bunodont teeth of hippos show a relatively

thick enamel layer that matches the anatomy of BL5-0 tooth fragment. Therefore, although the specimen has no clear anatomical resolution, the more parsimonious explanation (according to Occam's razor) is that BL5-0 is probably part of a hippo tooth.

Thus, tooth specimen BL02-J54-100 from the level D of Barranco León is the first human remain found at the Orce sites.

Paleoanthropology

The human specimen, BL02-J54-100 (Fig. 2), is a complete crown of an isolated lower left first deciduous molar (dm₁). It is heavily worn on the occlusal surface, category 5 according to Molnar (1971). Roots are missing due to resorption, suggesting that the tooth was shed ante-mortem. A distal wear facet is present at the contact with dm₂. The occlusal outline is oval and slightly asymmetrical. It presents the five main cusps. The protoconid is the largest and it is mesially displaced, followed by the metaconid, the hypoconid and finally the entoconid. The distal bifurcation of the central groove sets the limits of a small hypoconulid. At the dentine and pulp cavity level, the hypoconulid is only hinted as a faint elevation (Fig. 2). A well developed lingual groove and less developed buccal grooves are present. Mesial cusps represent the major part of the crown, and they are connected by a mid-trigonid crest. The tooth exhibits a strong mesial marginal ridge, prolonged by a vestigial paraconid, separated from the metaconid by a deep V-shaped groove that opens toward the lingual face. This feature is present in *Australopithecus* and *Homo*, such as KNM-ER 820, KNM-ER 1507 and the Zhoukoudian specimens (Weidenreich, 1937; Wood, 1991), and absent in *Paranthropus*. BL02-J54-100 exhibits a tuberculum molare in the mesiolingual angle, expanding onto the root trunk, but less developed than that found in the dm₁ from Gran Dolina-TD6, Atapuerca (Bermúdez de Castro et al., 1999) (Fig. S1). The presence of the tuberculum molare, the relative expansion of the mesial cusps and mesial marginal ridge, and the enlargement and relative position of the protoconid are classical diagnostic features in the genus *Homo* (Hillson, 1996; Keyser et al., 2000) (Fig. S1).

Overall, BL02 is a large tooth although there is a great overlap with the dm₁ dimensions of other hominin groups (Table 1 and Fig. S2). The mesiodistal (MD) diameter is close to the mean of *Paranthropus robustus* and *Australopithecus afarensis* but it also falls within the range of variation of African, Asian and European early Pleistocene *Homo* as well as *Homo heidelbergensis* and the Middle Paleolithic *Homo sapiens* sample. The buccolingual (BL) diameter is also close to the mean of *Paranthropus robustus* and *Australopithecus afarensis* and larger than that of any of the studied groups except for *Paranthropus boisei* and *Homo antecessor*. The crown computed area falls within the range of variation of *Paranthropus robustus*, *Australopithecus afarensis* and *H. heidelbergensis*. The highest values of the *Homo* sample are represented by the European early and middle Pleistocene fossils such as Arago 66, ATD6-94 and Barranco León together with the Neanderthal specimens from Teshik-Tash and Shanidar VII (Fig. S2).

Archeology

The lithic assemblage at Barranco León is composed of 1244 artefacts (Fajardo, 2008), including 26 cores, 185 whole flakes, 78 flake fragments, 759 waste flakes or débris, 17 retouched pieces, 92 angular fragments, 12 modified cobbles (including hammerstones) and 75 unmodified materials (cobbles and stones).

Flint, limestone and quartzite compose the raw material of the lithic assemblage. Jurassic formations situated around 3 km south of the site, contain several primary sources of good raw material, while several flint sources in secondary deposits of alluvial and colluvial origin were also found near the site (Toro-Moyano et al., 2009, 2011).




Figure 1. Barranco León site. Left: Geological and geographical location. Right: Section with paleomagnetic results and ESR ages in the middle.

Humans knapped at the Barranco León D site. Two sets of refitted flint pieces were identified. The first set is composed of four pieces: one core and three flakes (see Fig. 3a). Although it is not complete, the first and the last stages of the reduction sequence are

present. The first stage shows the application of the orthogonal technique with no preparation of the striking platform; cortical surfaces and previous negative scars were used as striking platforms. The last stage is composed of the exhausted nucleus with




Figure 2. Upper: Tooth specimen BL02-J54-100, left dm_1 from Barranco León D. a: Occlusal view; b: Buccal view; c: Distal view; d: Lingual view; and e: Mesial view. Lower: Computed tomographic reconstruction of the enamel and dentine surfaces (left and center) and the pulp cavity (right) of the dm_1 . The arrow points to the presence of a small hypoconulid.

ventral face exploitation of a flake showing the economy of good quality raw materials. The second refitting set is formed by pieces involuntarily detached during the knapping process.

Two main techniques were used for knapping, hard-hammer percussion and direct unipolar and bipolar technique (axial and non axial using an anvil). The choice of these techniques is related to the texture and quality of the raw material. The cobbles and tabular fragments of flint were destined for the production of flake cutting edges, while the less frequent limestone artefacts include cores, battered percutors and debitage.

The primary goal of the Barranco León knappers seems to have been obtaining small flakes, perhaps to fulfill immediate needs including rapidly cutting meat from large herbivore carcasses. The proportion of flakes, together with the high frequencies of débris (61.01%), broken and whole flakes, and angular fragments (28.54%), which sum up to 89.55% of debitage elements in the whole assemblage, and the refitting pieces show that the debitage was in situ. Although all stages of core reduction are presented, the most abundant are from final stages of flaking. The proportion of retouched artefacts (1.37%) is low and does not show stylistic standardization. These are basically light-duty retouched pieces (see Fig. 3b).

The striation marks and polished areas (Toro-Moyano et al., 2003) on the lithic material (Fig. 3c) are similar to those reported in well known African and European assemblages such as Olduvai Gorge, Koobi Fora or Monte Poggiolo (Keely and Toth, 1981; Sussman, 1987; Peretto et al., 1998). These marks suggest that the tools were used on a variety of materials.

The lithic artefacts are found together with large mammal remains showing evidences of anthropic activity, including true spiral or helical fractures, impact points, flake scars and bone flakes (Binford, 1981; Shipman, 1981a, b; Johnson, 1985; Blumenschine and Selvaggio, 1988) (Fig. 3d–i).

Chronology

Electron spin resonance (ESR) dating was applied to optically bleached quartz grains extracted from sediments. This method is based on the detection of paramagnetic centres, e.g., Aluminum (Al) or Titanium (Ti), that are created by the interaction of the natural radioactivity with the quartz sample (Ikeya, 1993). Similar to Optically Stimulated Luminescence (OSL) dating, ESR is an optical dating method that relies on the zeroing of any previously present ESR signal by sunlight exposure at the time of deposition (see details in Yokoyama et al., 1985). However, while OSL is usually limited to late middle Pleistocene time range, ESR may potentially go further back in time and cover the whole Quaternary time range (e.g., Voinchet et al., 2010), given the long term thermal stability of the ESR signal of the Al center (Toyoda and Ikeya, 1991) and its absence of saturation at high irradiation doses (Lin et al., 2006).

Five sediment samples for ESR dating (ESR BL-1 to ESR BL-5) were collected in situ in 2004, 2005 and 2006 from the sedimentary sequence of the excavated area at BL (Fig. 1). This study focused on Al signal because no measurable Ti signal was detected in our samples. Experimental conditions and age calculations are derived from Duval (2008) and detailed in the Supplementary Online Information.

Electron spin resonance age estimates are overall consistent with the stratigraphy, i.e., following a general increase with depth: they range from 1.02 ± 0.09 Ma to 1.88 ± 0.19 Ma and are all in agreement for attributing an early Pleistocene age to the deposits (Table S2). ESR BL-1 was sampled at the base of the sequence, about 1.5 m under the archaeological level D1, and yields a maximum age of 1.73 ± 0.17 Ma for the deposits. ESR BL-2 comes from D1 layer and provides an age of 1.46 ± 0.17 Ma. ESR BL-3 and ESR BL-4 samples were collected from level D2, which overlies D1: their ESR ages are 1.88 ± 0.19 Ma and 1.23 ± 0.12 Ma, respectively.

Table 1
Metric data of the specimen BL02-J54-100 from Barranco León, compared with other hominin samples.

Specimens	n	MD		BL		BL*MD (computed crown area)		BL/MD*100 (crown shape index)		Data source
		X	s.d.	X	s.d.	X	s.d.	X	s.d.	
<i>Paranthropus robustus</i> DNH 44, DNH 47, DNH 60, Kromdraai, SK61 (right), SK3978, SK3979	7	10.28	0.66	8.38	0.63	86.55	11.70	81.53	3.15	Robinson, 1956; Day, 1986; Keyser et al., 2000
<i>Paranthropus boisei</i> KNM-ER 1477, L704-2	2	11.65	–	9.30	–	108.41	–	80.03	–	Howell and Coppens, 1973; Wood, 1991
<i>Australopithecus africanus</i> Stw104, Stw151, Stw296, Sts24, Taung	5	8.77	0.32	7.50	0.45	65.90	5.82	85.53	3.98	Robinson, 1956; Moggi-Cecchi et al., 2006
<i>Australopithecus afarensis</i> AL333-43 (left), AL333-30, LH3, LH2	4	9.77	1.48	8.35	0.77	82.41	20.44	86.030	6.41	White, 1980; Johanson et al., 1982
Early <i>Homo</i> DNH 35	1	9.10	–	7.50	–	68.25	–	82.42	–	Keyser et al., 2000
<i>Homo erectus</i> African <i>Homo erectus</i> : KNM-ER 820 (right), KNM-ER 1507 Asian <i>Homo erectus</i> : Sinanthropus 123, 125	2 2	8.95 8.75	– –	7.30 6.80	– –	65.41 59.71	– –	81.49 78.57	– –	Leakey and Wood, 1973,1974 Weidenreich, 1937
<i>Homo antecessor</i> Gran Dolina-TD6 (ATDG-94) ^a	1	9.00	–	8.80	–	79.20	–	97.78	–	
<i>Homo heidelbergensis</i> Arago 34, 55, 66 ^a	3	9.43	0.66	8.03	0.40	75.96	9.30	85.23	1.71	
<i>Homo neanderthalensis</i> Archi, Gibraltar II, Kulna III, Pech de l'Azé, ^a Shanidar III, Shanidar VII, Chateaufeuf I, Chateaufeuf II, Krapina 67, Krapina A, Arcy 18, Arcy 25, Arcy 33, Roc de Marsal, Dederiyeh, Kebara, Teshik-Tash, Combe-Grenal, Taubach, La Chaise H13, La Chaise H14, La Chaise Bourgeois Delauney, Engis, La Ferrassie 8 ^a	24	8.83	0.44	7.56	0.47	66.86	6.23	85.68	5.46	Tillier, 1979; Wolpoff, 1979; Madre-Dupouy, 1985; Bailey and Hublin, 2006
<i>Homo sapiens</i> Middle Paleolithic: Skhül I, Skhül X, Qafzeh 4 Upper Paleolithic: Miesslingtal, Isturitz 72, Badger's Hole, Bruniquel 537, Bruniquel 540, Isturitz 4, Isturitz 9, Lagerie Basse 905, Lagerie Basse 909, La Madeleine 4, Le Figuer ^a	3 11	9.43 8.13	0.60 0.75	7.50 7.23	0.30 0.76	70.63 59.14	1.76 10.11	79.86 89.22	8.35 8.35	Tillier, 1979 Fraye, 1978
Contemporary: Musée de l'Homme collections (various geographical origins) ^a	21	8.31	0.36	6.96	0.26	57.99	3.95	83.88	3.79	
BL02-J54-100	1	10.05	–	8.48	–	85.22	–	84.4	–	

^a Measurement taken by authors.




Figure 3. Archaeological evidences from level D of Barranco León. a: Refitted pieces; b: Retouched piece; c: Striation and one polished marks; d–f: Bone cut marks, BL03-L62-2, rib fragment of megaherbivore (cf. *Hippopotamus antiquus*), it shows a long and curved cut mark oblique to the major edge of the rib in the central area of the bone, which is related to the evisceration process; g: Bone impact point; h–i: Bone flakes. Note: All lithic and fossil specimens shown in this figure were found during the excavations at Barranco León by the members of the research team that coauthor this article, with the only exception of specimen BL95 C1 A3 n.15, the central piece of the refitting group of Figure 3A, which was published in Gibert et al. (1998b).

Sample ESR BL-5, originating from level E1 at the top of the local sequence, yields a minimum age of 1.02 ± 0.09 Ma for the deposits.

Electron spin resonance age results obtained for layer D show some apparent scatter. More specifically, the age of ESR BL-3 looks somewhat overestimated in comparison with the other ages. This is probably due to the fitting of the dose response curve, which goes above the natural point and thus might induce an overestimation of the equivalent dose value (Fig. S3). Nevertheless, this age may not be discarded, regarding the small sample size ($n = 3$). Consequently, based on the three ESR samples BL-2 to BL-4, a weighted mean ESR age of 1.43 ± 0.38 Ma may be calculated for the layer D that

encloses the archaeological layer at Barranco León. The quite large final error may be interpreted as the consequence of the age scattering and the limited number of samples. This age is in quite good agreement with the ESR chronologies obtained for the nearby sites of Fuente Nueva-3 and Venta Micena, of 1.19 ± 0.21 Ma and ~ 1.4 Ma, respectively (Duval et al., 2011, 2012). However, one should be cautious in their interpretation, since the quite large error range does not allow any chronological distinction between the three sites from the Orce area. These results show the potential of ESR dating of quartz grains from early Pleistocene fluvio-lacustrine deposits. Future work will definitely need to be focused

on the improvement of the precision of this promising preliminary ESR chronological framework based on quartz grains extracted from sediment.

A new paleomagnetic study (Table S1), combined with previous results from the same stratigraphic section (Oms et al., 2000b) shows that the entire stratigraphic section of Barranco León has reverse polarity (Fig. 1). This fully reversed time interval for the whole silty calcareous member has been also confirmed in other sections by Scott et al. (2007) [see also Gibert et al. (2006) and comments in Agustí et al. (2007) and Oms et al. (2011)]. Consequently, the deposits may be correlated to the Matuyama chron (0.78–2.58 Ma; Gradstein et al., 2005), and more likely between Olduvai and Jaramillo subchrons, as indicated by the ESR chronology and faunal evidence.

Both micro and macrofauna support the ESR ages and the magnetostratigraphic interpretation. The combination of rodent species at the site, including *Mimomys savini*, *Allophaiomys* aff. *lavocati*, *Castillomys crusafonti* and *Apodemus* aff. *mystacinus*, further constrains its age. An age younger than the Olduvai subchron (1.95–1.77 Ma) is inferred from the more derived morphology of *Allophaiomys* aff. *lavocati* compared with *A. cf. deucalion* from the site of Kryzhanovka (Tesakov, 1998), which is associated with the Olduvai subchron (Pevner et al., 1998). *Allophaiomys* aff. *lavocati* is, in turn, more archaic than the microtine species present at Vallonet (France) and Untermassfeld (Germany), two localities dated to the Jaramillo subchron (0.99–1.07 Ma; Yokoyama et al., 1988; Wiegank, 1997). Therefore, Barranco León is placed within the reverse interval between the Jaramillo (0.99–1.07 Ma) and Olduvai (1.95–1.77 Ma) subchrons.

In the Guadix-Baza Basin, a biozonation based on rodents includes four biozones within this interval: *Mimomys oswaldoreigi* Zone, *Allophaiomys ruffoi* Zone, *Allophaiomys* aff. *lavocati* Zone and *Iberomys huescarensis* Zone (Agustí et al., 2010; Oms et al., 2011). The average duration of each biozone is 200 thousand years. The *Mimomys oswaldoreigi* Zone includes the sites of Barranco Conejos, Orce 2 and Orce D. The *Allophaiomys ruffoi* Zone includes the famous site of Venta Micena and other levels such as Cañada de Murcia 1, Fuente Nueva 2, Orce 4 and Orce 7 (Agustí et al., 2010). The *Allophaiomys* aff. *lavocati* Zone includes the sites of Barranco León D and Fuente Nueva 3. The *Iberomys huescarensis* Zone includes the sites of Huéscar 1, Puerto Lobo and Loma Quemada (Mazo et al., 1985; Agustí et al., 2010). The Last Occurrence (LO) of *Allophaiomys* aff. *lavocati* can be established below the base of the *Iberomys huescarensis* Zone, dated to the Jaramillo subchron at 1.07 Ma. The First Occurrence (FO) of *Allophaiomys* aff. *lavocati* is more difficult to establish, since there is no section in the basin where the boundary between the *Allophaiomys ruffoi* Zone and the *Allophaiomys* aff. *lavocati* Zone has been recognized. However, a recent dating of the level TE9 from Sima del Elefante, in the Atapuerca karstic complex (Cuenca-Bescós et al., 2001), contains a more evolved *Allophaiomys* than *A. aff. lavocati* from Barranco León D. The age of this level has been established at 1.22 ± 0.16 Ma, based on cosmogenic nuclides (Carbonell et al., 2008). Therefore, the age of the level BL D can be constrained between 1.77 Ma (top of Olduvai subchron) and ~ 1.2 Ma (age of Sima del Elefante).

However, interpolation of metric parameters measured on the lower first molar of diverse well dated early-middle Pleistocene microtine species provides a way to further constrain the age of the findings. The relative length of the anteroconid complex in microtines (the A/L parameter) (Meulen, 1973) has proven to be a useful tool for dating Plio-Pleistocene sites (Maul et al., 1998). The A/L values of *Allophaiomys* aff. *lavocati* from level BL D suggest an age older than Sima del Elefante and close to 1.4 Ma (Table S3).

The large mammals assemblage also points to an older chronology for BL-D than level TE9 from Sima del Elefante. In addition,

although it is commonly accepted that due to taphonomic reasons the lack of record of a taxon in an assemblage is not an evidence of its absence from the original community, the nonappearance of suids in any of the Late Villafranchian Orce sites is a strong argument that should not be avoided as a reliable biochronological marker. Suids are found in Europe before and during the Olduvai subchron, including Fonelas P-1 site in the Guadix-Baza Basin, dated to 2.0 Ma, which suid remains have been ascribed to *Potamochoerus magnus* (Arribas et al., 2009), as well as in many other localities with the presence of *Sus strozzi* (see Rook and Martínez-Navarro, 2010). However, there are no pigs in the chronological range comprised between the post Tasso Faunal Unit, in the base of the Late Villafranchian (~ 1.8 Ma), and their arrival in Western Europe at the level TE9 from Sima del Elefante (~ 1.2 Ma), where pigs are recorded under the name of *Sus* sp. (Carbonell et al., 2008), and at the site of Untermassfeld (Germany), dated 1.0–1.1 Ma and ascribed to *Sus scrofa prisca* (Güerin and Faure, 1997). At sites like Dmanisi (Georgia), as well as at Pirro Nord (Italy), Apollonia-1 (Greece), Sainzelles (France) and, of course, at the Orce sites, no record of pigs has never been documented. After 30 years of excavation of 350 m² at Venta Micena, 140 m² at Barranco León, and 106 m² at Fuente Nueva-3, and continuous research in the triangle formed by the region of Orce-Fuente Nueva-Venta Micena, pig remains have never been found among the more than 25 000 large mammal fossils unearthed from these sites, and this is probably the best record of the early Pleistocene fauna in Europe. When pigs are in the ecosystem, they use to be abundant in the large mammal community given their opportunistic feeding behavior and reproductive success. For this reason, they are usually well represented in the fossil assemblages. Suids are well recorded in the African and Asian archaeological and paleontological sites, including those from the Levantine Corridor as 'Ubeidiya (Geraads et al., 1986), or Evron Quarry (Tchernov et al., 1994)', but they are not definitively present in Orce or in any other sites of the European continent during the period comprised between 1.7 and 1.2 Ma. Later, the genus *Sus* is recorded everywhere in Europe. For this reason, the absence of suids from level BL D constrains the upper limit of the chronological range estimated for this site to >1.2 Ma, in agreement with the age estimated from the small mammal assemblage.

Conclusions

The specimen BL02-J54-100 found at Barranco León site, a human tooth with an estimated age by ESR between 1.02 and 1.73 Ma, and by paleomagnetism between 1.07 and 1.77 Ma, but close to 1.4 Ma according to the biochronologic evidence, represents the oldest anatomical evidence of human presence in Western Europe. This finding, combined with the important lithic tool assemblage from the level D of Barranco León, confirms that Western Europe was colonized soon after the first expansion out of Africa, currently documented at the Dmanisi site.

Acknowledgments

We thank all the members of the Orce research project involved in the excavations and the study of the Barranco León paleoanthropological site. We thank Professor P. Rivas for his support and encouragement. Also we thank Professor H. de Lumley and his team for their help. M.T. Alberdi, M.R. Palombo, L. Abbazzi, F. Lacomat, S. Bailón, H.A. Blain, J. Madurell, and M. Furió helped in the determination of the vertebrate fauna. C.F. and M.D. thank P. Voinchet and J. Despriée for their help during the ESR sampling. R. Grün and J.-J. Bahain provided helpful comments on the ESR dating. We thank Ph. Mennecier, A. Fort and V. Laborde from the MNHN (Musée de l'Homme) for the access to fossil and extant collections.

Photographs and microscan of the hominin molar were carried at CENIEH of Burgos, Spain. We thank L. Martín-Francés and M. Martínez de Pinillos for their help with the micro-CT analysis. ESR data were obtained at the Prehistory Department of MNHN (Paris, France). Paleomagnetic measurements were carried out at the SCT of the Barcelona University. Scanning photographs of the lithic artefacts were carried out at the CERP of Tautavel (France) by Mme. B. Deniaux. We thank Ms. Sheila Wglie for reviewing the English version of this paper. D. Begun, E. Delson, E. Carbonell, T. D. White, D. Lordkipanidze, S.C. Antón, M.M. Lahr, C. Stringer, M. Tappen, L. Rook, R. Ferring, the Associate Editor and three anonymous reviewers revised a first draft of the manuscript. This study has been sponsored by the Dirección General de Bienes Culturales (Consejería de Cultura, Junta de Andalucía, Spain), and it has been supported by projects CGL2010-15326, CGL2010-16821, CGL2012-38358, and CGL2008-04896 of the Spanish Ministry of Economy and Competitiveness.

Appendix A. Supplementary data

Supplementary data related to this article can be found online at <http://dx.doi.org/10.1016/j.jhevol.2013.01.012>.

References

- Agustí, J., Madurell, J., 2003. Los arvicólidos (Muroidea, Rodentia, Mammalia) del Pleistoceno inferior de Barranco León y Fuente Nueva 3 (Orce, Granada). Datos preliminares. In: Toro, I., Agustí, J., Martínez-Navarro, B. (Eds.), *El Pleistoceno Inferior de Barranco León y Fuente Nueva 3, Orce (Granada)*. E.P.G. Arqueología Monografías, vol. 17. Junta de Andalucía, Consejería de Cultura, pp. 137–145. Memoria Científica Campañas 1999–2002.
- Agustí, J., Oms, O., Parés, J.M., 2007. Biostratigraphy, paleomagnetism and geology of the Orce ravine (Southern Spain). Comment on the paper by Gibert et al. (2006). *Quatern. Sci. Rev.* 26, 568–572.
- Agustí, J., Blain, H.-A., Furió, M., De Marfá, R., Santos-Cubedo, A., 2010. The early Pleistocene small vertebrate succession from the Orce region (Guadix-Baza Basin, SE Spain) and its bearing on the first human occupation of Europe. *Quatern. Int.* 223–224, 162–169.
- Anadón, P., Gabàs, M., 2009. Palaeoenvironmental evolution of the early Pleistocene lacustrine sequence at Barranco León archaeological site (Orce, Baza Basin, Southern Spain) from stable isotopes and Sr and Mg chemistry of ostracod shells. *J. Paleolimnol.* 42, 261–279.
- Anadón, P., Utrilla, R., Julià, R., 1994. Palaeoenvironmental reconstruction of a Pleistocene lacustrine sequence from faunal assemblages and ostracode shell geochemistry, Baza Basin, SE Spain. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 111, 191–205.
- Arribas, A., Palmqvist, P., 2002. The first human dispersal to Europe: remarks on the archaeological and palaeoanthropological record from Orce (Guadix-Baza basin, southeastern Spain). *Hum. Evol.* 1–2, 55–78.
- Arribas, A., Garrido, C., Viseras, C., Soria, J.M., Pla, S., Solano, J.G., Garcés, M., Beamud, E., Carrión, J.S., 2009. A mammalian lost world in southwest Europe during the Late Pliocene. *Plos One* 4, e7127.
- Arzarello, M., Marcolini, F., Pavia, G., Pavia, M., Petronio, C., Petrucci, M., Rook, L., Sardella, R., 2009. L'industrie lithique du site Pléistocène inférieur de Pirro Nord (Apricena, Italie du sud): une occupation humaine entre 1,3 et 1,7 Ma. *L'Anthropologie* 113, 47–58.
- Bailey, S., Hublin, J., 2006. Dental remains from the Grotte du Renne at Arcy-sur-Cure (Yonne). *J. Hum. Evol.* 50, 485–508.
- Binford, L.R., 1981. *Bones: Ancient Men and Modern Myths*. Academic Press, New York.
- Blumenshine, R.J., Selvaggio, M.M., 1988. Percussion marks on bone surface as a new diagnostic of hominid behaviour. *Nature* 333, 763–765.
- Bermúdez de Castro, J.M., Rosas, A., Nicolás, M.E., 1999. Dental remains from Atapuerca-TD6 (Gran Dolina site, Burgos, Spain). *J. Hum. Evol.* 37, 523–566.
- Campillo, D., Cuesta, M.M., García-Guixé, E., Chimenos, E., Devenat, L., Baxarias, J., 2006. An occipital crest in an infant cranium from the Roman necropolis of Francolí (Tarragona, Spain): implications to the interpretation of the Orce skull. *Rev. Esp. Antropol. Fis.* 26, 93–101.
- Carbonell, E., Bermúdez de Castro, J.M., Parés, J.M., Pérez-González, A., Cuenca-Bescós, G., Ollé, A., Mosquera, M., Hugué, R., Made, J.V.D., Rosas, A., Sala, R., Vallverdú, J., García, N., Granger, D.E., Martínón-Torres, M., Rodríguez, X.P., Stock, G.M., Vergés, J.M., Allué, E., Burjachs, F., Cáceres, I., Canals, A., Benito, A., Díez, C., Lozano, M., Mateos, A., Navazo, M., Rodríguez, J., Rosell, J., Arsuaga, J.L., 2008. The first hominin of Europe. *Nature* 452, 465–469.
- Crochet, J.-Y., Welcomme, J.-L., Ivorra, J., Ruffet, G., Boulbes, N., Capdevila, R., Claude, J., Firmat, C., Métails, G., Michaux, J., Pickford, M., 2009. Une nouvelle faune de vertébrés continentaux, associée à des artefacts dans le Pléistocène inférieur de l'Hérault (Sud de la France), vers 1,57 Ma. *C.R. Palevol.* 8, 725–736.
- Cuenca-Bescós, G., Canudo, J.L., Laplana, C., 2001. La séquence des rongeurs (Mammalia) des sites du Pléistocène inférieur et moyen d'Atapuerca (Burgos, Espagne). *L'Anthropologie* 105, 115–130.
- Day, M.H., 1986. *Guide to Fossil Man*, fourth ed. The University of Chicago Press, Chicago.
- Despriée, J., Voinchet, P., Tissoux, H., Moncel, M.-H., Arzarello, M., Robin, S., Bahain, J.-J., Falguères, C., Courcimault, G., Dépont, J., Gageonnet, R., Marquer, L., Messager, E., Abdessadok, S., Puaud, S., 2010. Lower and middle Pleistocene human settlements in the middle Loire river basin, centre region, France. *Quatern. Int.* 223–224, 345–359.
- Duval, M., 2008. Evaluation du potentiel de la méthode de datation par Résonance de Spin Electronique (ESR) appliquée aux gisements du Pléistocène inférieur: étude des gisements d'Orce (bassin de Guadix-Baza, Espagne) et contribution à la connaissance des premiers peuplements de l'Europe. Ph.D. Dissertation. Muséum National d'Histoire Naturelle, Paris.
- Duval, M., Falguères, C., Bahain, J.-J., Grün, R., Shao, Q., Aubert, M., Hellstrom, J., Dolo, J.-M., Agustí, J., Martínez-Navarro, B., Palmqvist, P., Toro-Moyano, I., 2011. The challenge of dating early Pleistocene fossil teeth by the combined uranium series–electron spin resonance method: the Venta Micena palaeontological site. *J. Quatern. Sci.* 26, 603–615.
- Duval, M., Falguères, C., Bahain, J.-J., Grün, R., Shao, Q., Aubert, M., Dolo, J.-M., Agustí, J., Martínez-Navarro, B., Palmqvist, P., Toro-Moyano, I., 2012. On the limits of using combined U-series/ESR method to date fossil teeth from two Early Pleistocene archaeological sites of the Orce area (Guadix-Baza basin, Spain). *Quatern. Res.* 77, 481–482.
- Espigares, M.P., Martínez-Navarro, B., Palmqvist, P., Ros-Montoya, S., Toro, I., Agustí, J., Sala, R., 2013. *Homo vs. Pachyrocata*: earliest evidence of competition for an elephant carcass between scavengers at Fuente Nueva-3 (Orce, Spain). *Quatern. Int.* 295, 113–125.
- Fajardo, B., 2008. Les industries lithiques anciennes d'Orce: les sites de Barranco León et Fuente Nueva 3. Leur place dans le contexte des plus anciennes industries eurasiatiques. Ph.D. Dissertation. University of Montpellier.
- Fernández, J., Soria, J., Viseras, C., 1996. Stratigraphic architecture of the Neogene basins in the central sector of the Betic Cordillera (Spain): tectonic control and base level changes. In: Friend, P.F., Dabrio, J.C. (Eds.), *Record of Crustal Kinematics*. Cambridge University Press, Cambridge, pp. 353–364.
- Ferring, R., Oms, O., Agustí, J., Berna, F., Nioradze, M., Shelia, T., Tappen, M., Vekua, A., Zhvanida, D., Lordkipanidze, D., 2011. Earliest human occupations at Dmanisi (Georgian Caucasus) dated to 1.85–1.78 Ma. *Proc. Natl. Acad. Sci.* 108, 10432–10436.
- Frazer, D.W., 1978. *Evolution of the Dentition in Upper Paleolithic and Mesolithic Europe*. University of Kansas Press, Lawrence.
- Furió-Bruno, M., 2003. Los insectívoros (Mammalia) del Pleistoceno inferior de Fuente Nueva-3 y Barranco León (Orce, Granada). In: Toro, I., Agustí, J., Martínez-Navarro, B. (Eds.), *Geología, Paleontología, Paleocología Y Arqueología de la Depresión Guadix-Baza durante el Plio-Pleistoceno*. Arqueología Monografías, vol. 17. Consejería de Cultura, Junta de Andalucía, pp. 147–158.
- Geraads, D., Guérin, C., Faure, M., 1986. Les Suidés du Pléistocène ancien d'Oubeidiyeh (Israël). In: Tchernov, E. (Ed.), *Les Mammifères du Pléistocène Inférieur de la Vallée du Jordain a Oubeidiyeh*. Mémoires et Travaux du Centre de Recherche Français de Jérusalem, vol. 5. Association Paleorient, Paris, pp. 93–105.
- Gibert, J., Agustí, J., Moyà-Solà, S., 1983. Presencia de *Homo* sp. en el yacimiento del Pleistoceno inferior de Venta Micena (Orce, Granada). *Paleontol. Evol.* 1–12. *Publicación Especial*.
- Gibert, J., Campillo, D., Arqués, J.M., García-Olivares, E., Borja, C., Lowenstein, J.M., 1998a. Hominid status of the Orce cranial fragment reasserted. *J. Hum. Evol.* 34, 203–217.
- Gibert, J., Gibert, I., Iglesias, A., Maestro, E., 1998b. Two 'Oldowan' assemblages in the Plio-Pleistocene deposits of the Orce región, southeast Spain. *Antiquity* 72, 17–25.
- Gibert, J., Gibert, L.L., Albadalejo, S., Ribot, F., Sánchez, F., Gibert, P., 1999. Molar tooth fragment BL5-O: the oldest human remain found in the Plio-Pleistocene of Orce (Granada province, Spain). *Hum. Evol.* 14, 3–19.
- Gibert, J., Sánchez, F., Ribot, F., Gibert, L., Ferrandez, C., Iglesias, A., Gibert, P., 2002. Restes humaines dans les sédiments du pléistocène inférieur de la région d'Orce et de Cueva Victoria (au sud-est de l'Espagne). *L'Anthropologie* 106, 669–683.
- Gibert, L., Scott, G., Ferrández-Cañadell, C., 2006. Evaluation of the Olduvai subchron in the Orce ravine (SE Spain). Implications for the Plio-Pleistocene mammal biostratigraphy and the age of the Orce archaeological sites. *Quatern. Sci. Rev.* 25, 507–525.
- Gradstein, F.M., Ogg, J.G., Smith, A.G., 2005. *A Geologic Time Scale 2004*. Cambridge University Press, Cambridge.
- Guérin, C., Faure, M., 1997. The wild boar (*Sus scrofa priscus*) from the post-Villafranchian lower Pleistocene of Untermassfeld. In: Kahlke, R.-D. (Ed.), *Das Pleistozän von Untermassfeld bei Meningen (Thüringen)*. Romisch-Germanisches Zentralmuseum, Bonn, pp. 375–383.
- Hillson, S., 1996. *Dental Anthropology*. Cambridge University Press, Cambridge.
- Howell, F.C., Coppens, Y., 1973. Deciduous teeth of Hominidae from Pliocene-Pleistocene of lower Omo basin. *J. Hum. Evol.* 2, 461–472.
- Ikeya, M., 1993. *New Applications of Electron Spin Resonance Dating, Dosimetry and Microscopy*. World Scientific, Singapore.
- Johanson, D.C., White, T.D., Coppens, Y., 1982. Dental remains from the Hadar Formation, Ethiopia: 1974–1977 collections. *Am. J. Phys. Anthropol.* 57, 545–603.

- Johnson, E., 1985. Current developments in bone technology. In: Lemoine, G.M., MacEachern, A.S. (Eds.), 1985. *Advances in Archaeological Method and Theory*, vol. 8, pp. 157–235.
- Keely, L.H., Toth, N., 1981. Microwear polishes on early stone tools from Koobi Fora, Kenya. *Nature* 293, 464–465.
- Keyser, A., Menter, C., Moggi-Cecchi, J., Pickering, T.R., Berger, L., 2000. Drimolen: a new hominid-bearing site in Gauteng, South Africa. *S. Afr. J. Sci.* 96, 193–197.
- Leakey, R.E., Wood, B.A., 1973. New evidence of the genus *Homo* from East Rudolf, Kenya. *Am. J. Phys. Anthropol.* 39, 355–368.
- Leakey, R.E., Wood, B.A., 1974. New evidence of the genus *Homo* from East Rudolf, Kenya (IV). *Am. J. Phys. Anthropol.* 41, 237–243.
- Lin, M., Yin, G., Ding, Y., Cui, Y., Chen, K., Wu, C., Xu, L., 2006. Reliability study on ESR dating of the aluminium center in quartz. *Radiat. Meas.* 41, 1045–1049.
- Lordkipanidze, D., Jashashvili, T., Vekua, A., Ponce de León, M.S., Zollikofer, C.P.E., Rightmire, G.P., Pontzer, H., Ferring, R., Oms, O., Tappen, M., Bukhsianidze, M., Agustí, J., Kahlke, R., Kiladze, G., Martínez-Navarro, B., Moushkelishvili, A., Nioradze, M., Rook, L., 2007. Postcranial evidence from early *Homo* from Dmanisi, Georgia. *Nature* 449, 305–310.
- Madre-Dupouy, M., 1985. Les dents déciduales de l'enfant néanderthalien du Roc de Marsal – Dordogne – France. Etude analytique et comparative. Deuxième partie: la dentition inférieure. *L'Anthropologie* 89, 93–109.
- Martínez-Navarro, B., 2002. The skull of Orce: parietal bones or frontal bones? *J. Hum. Evol.* 42, 265–270.
- Martínez-Navarro, B., Turq, A., Agustí, J., Oms, O., 1997. Fuente Nueva-3 (Orce, Granada, Spain) and the first human occupation of Europe. *J. Hum. Evol.* 33, 611–620.
- Martínez-Navarro, B., Palmqvist, P., Madurell-Malapeira, J., Ros-Montoya, S., Espigares, M.P., Torregrosa, V., Pérez-Claros, J.A., 2010. La fauna de grandes mamíferos de Fuente Nueva-3 y Barranco León-5: estado de la cuestión. In: Toro, I., Martínez-Navarro, B., Agustí, J. (Eds.), *Occupaciones Humanas en el Pleistoceno Inferior y Medio de la Cuenca de Guadix-Baza*. Junta de Andalucía, Consejería de Cultura, Sevilla, pp. 197–236.
- Martínez-Navarro, B., Toro, I., Agustí, J., 2005. The large mammals assemblages from Venta Micena, Fuente Nueva-3 and Barranco León-5 (Orce). Early Pleistocene faunal and human dispersals into Europe. In: Molines, N., Moncel, M.-H., Monnier, J.-L. (Eds.), *Proceedings of the Symposium Entitled 'Recent Advances Concerning Settlement Changes as Well as Chronostratigraphic, Geological and Paleogeographical Framework of the Industries of the Middle and Early Paleolithic Period in Europe'*, Rennes, 22–25 September, 2003. John & Erica Hedges Ltd., Oxford, pp. 125–133. *British Archaeological Reports International Series S1364*.
- Maul, L., Masini, F., Abbazzi, L., Turner, A., 1998. The use of different morphometric data for absolute age calibration of some South- and Middle European arvicolid populations. *Palaeontogr. Ital.* 85, 111–151.
- Mazo, A.V., Sesé, C., Ruiz-Bustos, A., Peña, J.A., 1985. Geología y paleontología de los yacimientos Plio-Pleistocenos de Huéscar (Depresión de Guadix-Baza, Granada). *Estud. Geol.* 41, 467–493.
- Meulen, A.v.d., 1973. Middle Pleistocene smaller mammals from the Monte Peglia (Orvieto, Italy) with special reference to the phylogeny of *Microtus* (Arvicolidae, Rodentia). *Quaternaria* 17, 1–144.
- Moggi-Cecchi, J., Grine, F.E., Tobias, P.V., 2006. Early hominid dental remains from Members 4 and 5 of the Sterkfontein formation (1966–1996 excavations): catalogue, individual associations, morphological descriptions and initial metric analysis. *J. Hum. Evol.* 50, 239–328.
- Molnar, S., 1971. Human tooth wear, tooth function and cultural variability. *Am. J. Phys. Anthropol.* 34, 175–190.
- Moyà-Solà, S., Köhler, M., 1997. The Orce skull: anatomy of a mistake. *J. Hum. Evol.* 33, 91–97.
- Oms, O., Agustí, J., Gabàs, M., Anadón, P., 2000a. Lithostratigraphical correlation of micromammal sites and biostratigraphy of the Upper Pliocene to Lower Pleistocene in the Northeast Guadix-Baza Basin (southern Spain). *J. Quatern. Sci.* 15, 43–50.
- Oms, O., Parés, J.M., Martínez-Navarro, B., Agustí, J., Toro, I., Martínez-Fernández, G., Turq, A., 2000b. Early human occupation of western Europe: Paleomagnetic dates for two paleolithic sites in Spain. *Proc. Natl. Acad. Sci.* 97, 10666–10670.
- Oms, O., Anadón, P., Agustí, J., Julià, R., 2011. Geology and chronology of the continental Pleistocene archeological and paleontological sites of the Orce area (Baza basin, Spain). *Quatern. Int.* 243, 33–43.
- Palmqvist, P., 1997. A critical re-evaluation of the evidence for the presence of hominids in lower Pleistocene times at Venta Micena, Southern Spain. *J. Hum. Evol.* 33, 83–89.
- Palmqvist, P., Martínez-Navarro, B., Toro, I., Espigares, M.P., Ros-Montoya, S., Torregrosa, V., Pérez-Claros, J.A., 2005. Réévaluation de la présence humaine au Pléistocène inférieur dans le Sud de l'Espagne. *L'Anthropologie* 109, 411–450.
- Peretto, C., Amore, F.O., Antoniazzi, Alb., Antoniazzi, Ald., Bahain, J.J., Cattani, L., Cavallini, E., Esposito, P., Falguères, C., Gagnepain, J., Hedley, I., Laurent, M., Lebreton, V., Longo, L., Milliken, S., Monegati, P., Ollé, A., Pugliese, N., Renault-Miskovsky, J., Sozzi, M., Ungaro, S., Vannucci, S., Verges, J.M., Wagner, J.J., Yokoyama, Y., 1998. L'industrie lithique de CÀbelvedere di Monte Poggiolo: stratigraphie, matière première, typologie, remontages et traces d'utilisation. *L'Anthropologie* 102, 343–465.
- Pevner, M., Tesakov, A., Vengengeim, E., 1998. The position of the Tidzar Locality (Taman Peninsula, Russia) in magnetochronological scale. *Paludicola* 2, 95–97.
- Robinson, J.T., 1956. Dentition of Australopithecinae. *Transvaal Museum, Pretoria*.
- Rook, L., Martínez-Navarro, B., 2010. Villafranchian: the long story of a Plio-Pleistocene European large mammal biochronologic unit. *Quatern. Int.* 219, 134–144.
- Sanz de Galdeano, C., Vera, J.A., 1992. Stratigraphic record and palaeogeographical context of the Neogene basins in the Betic Cordillera. *Basin Res.* 4, 21–36.
- Scott, G., Gibert, L., Gibert, J., 2007. Magnetostratigraphy of the Orce region (Baza basin), SE Spain: new chronologies for early Pleistocene faunas and hominid occupation sites. *Quatern. Sci. Rev.* 26, 415–435.
- Shipman, P., 1981a. Applications of scanning electron microscopy to taphonomic problems. In: Cantwell, A.M., Griffin, J.B., Rothschild, N.A. (Eds.), *The Research Potential of Anthropological Museum Collections*. Ann. N.Y. Acad. Sci., vol. 376, pp. 357–385.
- Shipman, P., 1981b. *Life History of a Fossil: an Introduction to Taphonomy and Paleoecology*. Harvard University Press, Cambridge.
- Sussman, C., 1987. Résultats d'une étude de microtraces d'usure sur un échantillon d'artefacts d'Olduvai (Tanzanie). *L'Anthropologie* 91, 375–380.
- Tchernov, E., Horwitz, L.K., Ronen, A., Lister, A., 1994. The faunal remains from Evron Quarry in relation to other Lower Paleolithic hominid sites in the Levant. *Quatern. Res.* 42, 328–339.
- Tesakov, A., 1998. Early stage of *Allophaiomys* in eastern Europe. *Paludicola* 2, 98–105.
- Tillier, A.M., 1979. Restes crâniens de l'enfant moustérien Homo 4 de Qafzeh (Israël). La mandibule et les maxillaires. *Paléorient* 5, 67–85.
- Toro-Moyano, I., de Lumley, H., Barsky, D., Celiberti, V., Cauche, D., Moncel, M.-H., Fajardo, B., Toro, M., 2003. Las industrias líticas de Barranco León y Fuente Nueva 3 de Orce. Estudio técnico y tipológico. Las cadenas operativas. Análisis traceológico. Resultados preliminares. In: Toro, I., Agustí, J., Martínez-Navarro, B. (Eds.), *El Pleistoceno Inferior de Barranco León y Fuente Nueva 3, Orce (Granada)*. Memoria Científica Campañas 1999–2002. Junta de Andalucía, Consejería de Cultura, Sevilla, pp. 183–206.
- Toro-Moyano, I., de Lumley, H., Fajardo, B., Barsky, D., Cauche, D., Celiberti, V., Gregoire, S., Martínez-Navarro, B., Espigares, M.P., Ros-Montoya, S., 2009. L'industrie lithique des gisements du Pléistocène inférieur de Barranco León et Fuente Nueva 3 à Orce, Grenade, Espagne. *L'Anthropologie* 113, 111–124.
- Toro-Moyano, I., Barsky, D., Cauche, D., Celiberti, V., Grégoire, S., Lebegue, F., Moncel, M.H., de Lumley, H., 2011. The archaic stone tool industry from Barranco León and Fuente Nueva 3 (Orce, Spain): evidence of the earliest hominin presence in southern Europe. *Quatern. Int.* 243, 80–91.
- Toyoda, S., Ikeya, M., 1991. Thermal stabilities of paramagnetic defect and impurity centers in quartz: basis for ESR dating of thermal history. *Geochem. J.* 25, 437–445.
- Turq, A., Martínez-Navarro, B., Palmqvist, P., Arribas, A., Agustí, J., Rodríguez-Vidal, J., 1996. Le Plio-Pleistocene de la région d'Orce, province de Grenade, Espagne: bilan et perspectives de recherche. *Paleo.* 8, 161–204.
- Vekua, A., Lordkipanidze, D., Rightmire, G.P., Agustí, J., Ferring, R., Maisuradze, G., Moushkelishvili, A., Nioradze, M., Ponce de León, M., Tappen, M., Tvalchrelidze, M., Zollikofer, C., 2002. A new skull of early *Homo* from Dmanisi, Georgia. *Science* 297, 85–89.
- Vera, J.A., 1970. Estudio estratigráfico de la Depresión de Guadix-Baza. *Bol. Geol. Min* 84, 429–462.
- Vera, J.A., Fernández, J., López-Garrido, A.C., Rodríguez-Fernández, J., 1985. Geología y estratigrafía de los materiales plioceno-pleistocenos del sector Orce-Venta Micena (prov. Granada). *Paleontol. Evol.* 18, 3–11.
- Voinchet, P., Despriée, J., Tissoux, H., Falguères, C., Bahain, J.J., Gageonnet, R., Dépont, J., Dolo, J.M., 2010. ESR chronology of alluvial deposits and first human settlements of the Middle Loire Basin (Region Centre, France). *Quatern. Geochronol.* 5, 381–384.
- Weidenreich, F., 1937. The dentition of *Sinanthropus pekinensis*: a comparative odontography of the hominid. *Palaeont. Sinica D* 1, 1–180.
- White, T.D., 1980. Additional fossil hominids from Laetoli, Tanzania: 1976–1979 specimens. *Am. J. Phys. Anthropol.* 53, 487–504.
- Wiegand, F., 1997. Paläomagnetische Charakteristik des Unterpleistozäns von Untermassfeld. In: Kahlke, R.D. (Ed.), *Das Plesitozän von Untermassfeld bei Meiningen (Thüringen)*. Romisch-Germanisches Zentralmuseum, Bonn, pp. 63–69.
- Wolpoff, M.H., 1979. The Krapina dental remains. *Am. J. Phys. Anthropol.* 50, 67–114.
- Wood, B.A., 1991. Koobi Fora Research Project. In: *Hominid Cranial Remains from Koobi Fora*, vol. 4. Clarendon Press, Oxford.
- Yokoyama, Y., Falguères, C., Quaegebeur, J.P., 1985. ESR dating of quartz from quaternary sediments: first attempt. *Nucl. Tracks Rad. Meas.* 10, 921–928.
- Yokoyama, Y., Bibron, R., Falguères, C., 1988. Datation absolue des planchers stalagmitiques de la grotte du Vallonnet à Roquebrune-Cap-Martin (Alpes-Maritimes) France, par la résonance de spin électronique (ESR). *L'Anthropologie* 92, 429–436.